


Yeast Extract Agar for Molds

Cat. 1312

For the cultivation of yeast and molds from diverse materials, especially milk and dairy products

Practical information

Applications	Categories
Growth	Yeasts and molds
Industry: Dairy products	



Principles and uses

Yeast Extract Agar for Molds is suitable to cultivate molds and yeasts from milk and dairy products.

Yeast extract is a source of vitamins, particularly of the B-group essential for bacterial growth. Dextrose is the carbohydrate energy source. Bacteriological agar is the solidifying agent. The high dextrose concentration makes this medium selective for yeasts and molds.

Formula in g/L

Bacteriological agar	20	Dextrose	10
Yeast extract	5		

Preparation

Suspend 35 grams of the medium in one liter of distilled water. Mix well and dissolve by heating with frequent agitation. Boil for one minute until complete dissolution. Sterilize in autoclave at 121°C for 15 minutes. Cool to 45-50°C, mix well and dispense into plates.

Instructions for use

The inoculation method can be carried out either by flooding or on the surface, depending on the purpose for which the medium is intended. Incubation time is 7 days at a temperature of 28°C and in an aerobic environment.

For increased selectivity, acidify the medium or add antibiotics with the aseptic technique. It is not recommended to heat the acidified agar medium.

Quality control

Solubility	Appearance	Color of the dehydrated medium	Color of the prepared medium	Final pH (25°C)
w/o rests	Fine powder	Beige	Amber, slightly opalescent	6,5 ± 0,2 a 25°C

Microbiological test

Incubation conditions: 28°C / 7 days

Microorganisms	Specification
Candida albicans ATCC 10231	Good growth

Aspergillus brasiliensis ATCC 16404
Escherichia coli ATCC 25922
Staphylococcus aureus ATCC 25923

Good growth
Good growth
Good growth

Storage

Temp. Min.:2 °C
Temp. Max.:25 °C

Bibliography

Cooke, W.B. and A. R. Brazis. 1968. Occurrence of molds and yeasts in dairy products. Mycopathol. Mycol. Appl. 35:281 -289. Overcase, W.W. and D.J. Weakley. 1969. An aureomycin-rose Bengal agar for enumeration of yeast and mold in cottage cheese. International Dairy Federation. Standard Method ISO/DIS 6611.

Koburger, J.A.. 1970. Fungi in foods: 1. Effect of inhibitor and incubation temperature on enumeration. J. Milk Food Technol. 33:433-434.