

Eosin Methylene Blue Agar (EMB)


Cat. 1039

For the isolation, cultivation and differentiation of Gram negative enteric bacilli from clinical and other specimens

Practical information

Applications	Categories
Selective isolation	Gram-negative enteric bacilli
Differentiation	Gram-negative enteric bacilli

Industry: Clinical



Principles and uses

Eosin Methylene Blue Agar (EMB) is a differential medium similar to Levine EMB Agar (Cat. 1050), used for the isolation of Enterobacteria. The use of eosin Y and methylene blue enable differentiation between lactose-fermenting and non-fermenting organisms. It is widely used in medical bacteriology, in techniques recommended by APHA and for the detection and enumeration of coliforms, contaminants of foods and drinking water.

Peptone provide nitrogen, vitamins, minerals and amino acids essential for growth. Sucrose is added to Lactose as a fermentable carbohydrate to detect coliforms that ferment sucrose more readily than lactose. Eosin Y and methylene blue dyes are both partial inhibitors of Gram-positive bacteria and pH indicators. Due to the lactose and sucrose, this medium can be differential in primary culture: Salmonellae and Shigellae which are lactose-negative can be differentiated from other lactose-negative and sucrose-positive organisms such as Proteus vulgaris, Citrobacter and Aeromonas. Dipotassium phosphate acts as a buffer system and bacteriological agar is the solidifying agent.

Formula in g/L

Bacteriological agar	13,5	Bacteriological peptone	10
Dipotassium phosphate	2	Eosin Y	0,4
Lactose	5	Methylene blue	0,065
Sucrose	5		

Typical formula g/L * Adjusted and/or supplemented as required to meet performance criteria.

Preparation

Suspend 36 grams of medium in one liter of distilled water. Mix well and dissolve by heating with frequent agitation. Boil for one minute until complete dissolution. Sterilize in autoclave at 121 °C for 15 minutes. Cool to 45-50 °C, mix well, avoiding the formation of bubbles and dispense carefully into Petri Dishes. DO NOT OVERHEAT.

Instructions for use

For the isolation of enteric pathogens from clinical samples. For clinical diagnosis, the type of sample is urine and feces.

- Inoculate a plate of EMB Agar and streak for isolation, allowing discrete colonies to develop.
- Incubate at 35±2 °C and observe at 24 hours and again at 48 hours.

Colonies characteristics:

- Salmonella and Shigella colonies are translucent and amber colored or colorless.
- Coliforms that use lactose and/or sucrose produce blue-black colonies with dark centers and a greenish metallic sheen.
- Other coliforms such as Enterobacter form mucoid, pink colonies.
- Strains of Enterococcus faecalis are partially inhibited on this medium and appear as colorless colonies.

Note: As the medium is moderately inhibitory some staphylococci, streptococci and yeast may grow. Also some Gram-negative non-fermenting bacilli may appear as non-lactose fermenters. Further Biochemical tests are necessary for genus identification.

Quality control

Solubility	Appearance	Color of the dehydrated medium	Color of the prepared medium	Final pH (25°C)
w/o rests	Fine powder	Purple-rose	Tournaol blue. After sterilization: Orange	7,2±0,2

Microbiological test

Incubation conditions: (35±2 °C / 24-48 h).

Microorganisms	Specification	Characteristic reaction
<i>Pseudomonas aeruginosa</i> ATCC 10145	Good growth	Colorless colony
<i>Klebsiella aerogenes</i> ATCC 13048	Good growth	Pink colony
<i>Salmonella typhimurium</i> ATCC 14028	Good growth	Colorless colony
<i>Escherichia coli</i> ATCC 25922	Good growth	Green colony with metallic shine
<i>Staphylococcus aureus</i> ATCC 25923	Inhibited growth	

Storage

Temp. Min.: 2 °C
Temp. Max.: 25 °C

Bibliography

American Public Health Association. Diagnostic Procedures and Reagents. 2nd Ed. APHA, Inc. New York, 1950
A.P.H.A Examination of dairy products. 10th Ed. APHA, Inc. New York, 1953. Society of American Bacteriologists. Manual of Microbiological Methods MacGraw-Hill New York, 1957.